

Journals Offprint Order Form



A member of the
Hodder Headline
Group

Arnold Journals
338 Euston Road
London NW1 3BH
Tel: +44 (0)20 7873 6000
Fax: +44 (0)20 7873 6325

This form should be returned at once to the above address

- Title of Journal: Palliative Medicine 16 (3)
- 1st Author:

A. FREE OFFPRINTS - 25 offprints of your article will be supplied free of charge

Please indicate opposite, the name and full postal address to whom they should be sent. In the case of multi-author articles, free offprints are only sent to the corresponding author.

B. PURCHASE OF ADDITIONAL OFFPRINTS

Please note that if an article is by more than one author, only one offprint form is sent and all offprints should be ordered on that form in consultation with the coauthors.

Offprint Price List (£ sterling UK and Europe; US\$ Rest of World)

	25		50		100		150		200	
	£	\$	£	\$	£	\$	£	\$	£	\$
1-4 pages	54	98	69	128	109	194	163	295	197	354
5-8 pages	72	130	95	171	145	259	199	357	259	466
9-16 pages	92	170	109	194	163	295	216	391	287	517
17-24 pages	109	194	124	224	182	324	257	463	327	588
Extra 8 pages	12	18	17	28	19	34	23	41	32	59

For larger quantities contact the publisher for a quotation.

Add 100% for any offprints including colour reproduction.

I wish to purchaseadditional offprints

ADDRESS FOR DELIVERY

(please print in capitals)

IMPORTANT

1. Cheques drawn on a UK or US bank should be made payable to Hodder Headline Group. We are unable to accept credit or debit card payments.
2. Orders will not normally be mailed until the publisher is in receipt of either the appropriate payment or an official purchase order.
3. The above are prepublication prices and apply only to orders received before the publication goes to press.
4. All despatches are by surface mail, normally within four weeks of publication.
5. Claims cannot be considered more than three months after despatch.

VAT will be added to UK invoices. Members of the EU will be required to pay VAT unless a VAT number is provided with order.

ADDRESS FOR INVOICE (please print in capitals)

Payment enclosed

Please invoice

Official order follows

Official order attached

no.

Signed.....

Date...../...../.....

Integration of acupuncture into the oncology clinic[†]

Peter AS Johnstone Radiation Oncology Service, Naval Medical Center, San Diego, California 92134-1014 and Radiation Oncology Division, University of California, San Diego, California 92103, **Gregory R Polston** Department of Anesthesiology, **Richard C Niemtzow** Radiation Oncology Service and Department of Anesthesiology and **Peter J Martin** Head and Neck Surgery Service, Naval Medical Center, San Diego, California 92134-1014

Abstract: Purpose/Objective: Patients with cancer or symptoms referable to cancer therapy were offered acupuncture as potential palliation of their symptoms. This paper describes the physical integration of the discipline into the Oncology Clinic, and patient perspectives on its availability and efficacy. **Patients and Methods:** Between August 1999 and May 2000, 123 patients with varying symptoms received acupuncture in our Center's Radiation and Medical Oncology Clinics and Breast Health Center. These patients had 823 visits during this time period. A practice outcome analysis was performed on patients receiving therapy between 1 January 2000 and 30 April 2000. The 89 patients treated during this interval had 444 total visits. In June and July 2000, a questionnaire was administered by phone to 79 of these patients (88.7%). Standard allopathic care continued while patients were receiving acupuncture. **Results:** Major reasons for referral included pain (53%), xerostomia (32%), hot flashes (6%) and nausea/loss of appetite (6%). Patients had a mean of five acupuncture visits (range 1–9). Most patients (60%) showed at least 30% improvement in their symptoms. About one-third of patients had no change in severity of symptoms. There were no untoward effects reported related to the acupuncture. When analysed by diagnosis, these values persist. Irrespective of response to therapy, 86% of respondents considered it 'very important' that we continue to provide acupuncture services. **Conclusion:** Acupuncture may contribute to fewer symptoms for cancer patients. Expansion of providers, continued patient follow-up, optimization of techniques and prospective objective measurement of response continue in our clinic. *Palliative Medicine* 2002; 16

Key words: acupuncture; alternative and complementary therapies; cancer; nausea; palliation; xerostomia

Resumé: Objectifs: nous avons proposé à des patients atteints de cancer ou de symptômes releves d'un traitement anti_cancéreux de l'acupuncture comme traitement palliatif de leur symptôme. Cet article décrit l'intégration physique de cette discipline à une activité d'oncologie et les points de vue du patient sur sa mise en œuvre et son efficacité. **Patient et méthode:** entre 8_99 et 5/00, 123 patients porteurs de divers symptômes ont reçu de l'acupuncture dans notre centre de traitement des pathologies mammaires et cliniques oncologiques médicales et radiothérapeutiques. Ces patients ont reçu 823 visites pendant cette période. Nous avons réalisé une analyse des résultats de la pratique recevant une thérapie entre le 1/1/00 et 4/30/00. Les quatre vingt neuf patients traités durant cet intervalle ont eu un total de 444 visites. En juin et juillet 2000, nous avons soumis un questionnaire par téléphone à 79 de ces patients (88,7%). Les soins allopathiques standards ont été poursuivis en même temps que l'acupuncture. **Résultats:** les principaux motifs de traitements étaient la douleur (53%), la xérostomie (32%), les bouffées de chaleur (6%) et les nausées/perde d'appétit (6%). Les patients ont eu une moyenne de 5 séances d'acupuncture (de 1 à 9). La plupart des patients (60%) ont montré une amélioration d'au moins 30% de leur symptôme. Dans environ 1/3 des patients, il n'y a pas eu de changement dans la sévérité des symptômes. Il n'y a pas eu d'effet indésirable lié à l'acupuncture. Ces résultats restent valables pathologies par pathologies. Quelque soit la réponse au traitement, 86% des participants considéraient que l'acupuncture était très importante de telle manière qu'elle a été

Address for correspondence: Dr Peter AS Johnstone, Radiation Oncology Division, Suite 14, Naval Medical Center, 34800 Bob Wilson Dr., San Diego, California 92134-1014, USA.

E-mail: pajohnstone@nmcscd.med.navy.mil

[†]The opinions and assertions contained herein are those of the authors and should not be construed as official or representing the views of the United States Navy or Department of Defense. Presented in part at the 83rd Annual Meeting of the American Radium Society, Coronado, CA, 29 April 2001.

poursuivie. **Conclusion:** l'acupuncture peut contribuer à la prise en charge de quelques symptômes chez des patients cancéreux. Dans notre clinique nous augmentons les praticiens, continuons à suivre les patients, optimisons les techniques et mesurons de façon objective et prospective les réponses. *Palliative Medicine* 2002; **16**

Mots-clés: acupuncture; thérapies alternatives et complémentaires; cancer; nausées; traitement palliatifs; xérostomie

Introduction

Acupuncture is gaining momentum and adherents as a valid intervention in palliative medicine. In the United States, this mirrors a movement towards complementary therapies; recent data reveal that 34–40%^{1,2} of patients seek nontraditional medical attention.

Acupuncture received needed validation at a National Institutes of Health (NIH) consensus conference in November of 1997. After review of the existing literature, the attendees concluded:

*Acupuncture as a therapeutic intervention is widely practised in the United States. While there have been many studies of its potential usefulness, many of these studies provide equivocal results because of design, sample size, and other factors. The issue is further complicated by inherent difficulties in the use of appropriate controls, such as placebos and sham acupuncture groups. However, promising results have emerged, for example, showing efficacy of acupuncture in adult post-operative and chemotherapy nausea and vomiting and post-operative dental pain. There are other situations such as addiction, stroke rehabilitation, headache, menstrual cramps, tennis elbow, fibromyalgia, myofascial pain, osteoarthritis, low back pain, carpal tunnel syndrome and asthma where acupuncture may be useful as an adjunct treatment or an acceptable alternative or be included in a comprehensive management programme. Further research is likely to uncover additional areas where acupuncture interventions will be useful.*³

Two recent literature reviews^{4,5} have shown its efficacy in pain, and called for further research in other arenas. Formal clinical trials of acupuncture have begun.⁶ Positive results are beginning to emerge in reputable, peer-reviewed journals.^{7–9}

In response to this movement, it was decided to integrate acupuncture into patient care at Naval Medical Center San Diego (NMCS D). In general, medical services and medications at NMCS D are offered to active-duty military personnel in the US Armed Forces and their dependents on a no-cost basis. Retirees and their dependents may gain access to NMCS D either on a space-available basis, or by enrolling in the US Department of Defense healthcare plan, TRICARE. There has never been a per-visit charge for any oncology or acupuncture services at NMCS D, and no patients were sent elsewhere because of inability to deal with their underlying oncology problem during this interval. Thus, access and ability to pay are two variables existing in the

US population at large that do not exist in the microcosm of the NMCS D population discussed here.

The first step in acupuncture integration was to add a Physician Acupuncturist to the medical staff. Although acupuncture may be administered by nonphysician providers in the United States, for accreditation purposes within the Department of Defense, a physician was preferable. Physician acupuncturists, like their nonphysician counterparts, are licensed at the State level; in almost all cases, licensure is automatic after completion of extensive training such as the 300 credit UCLA course 'Medical Acupuncture for Physicians'. The programme was initiated using a single acupuncturist (RCN) so trained, with over 10 years of experience.

For many reasons, including acupuncture's value in symptom palliation, the sites of initial integration were the Oncology Clinics. Patients with symptoms related to cancer or to cancer treatments were offered acupuncture as potential palliation of their symptoms. Although refractory pain and nausea were the two diagnoses expected to comprise most of the therapeutic interventions, early success in patients with radiation-induced xerostomia^{7,10} were an enormous and gratifying surprise.

After the programme had been in place for a year, additional providers were credentialed by the Medical Center, and nononcology patients were allowed access. However, prior to expansion of the service, a practice analysis was performed to document initial outcomes. This manuscript provides an overview of patient perspectives on this palliative programme, describes the quality assurance, peer review and oversight techniques currently in use, and lists current diagnoses being treated.

Patients and methods

In the first 9 months of the acupuncture integration, between August 1999 and May 2000, 123 patients with

- What did you have treated?
- Did you get better, worse, or stay the same?
- What percent change?
- Are you continuing or would you continue acupuncture?
- How important is it for our medical center to continue this form of therapy?

Figure 1 Questions asked via telephone

Table 1 Demographics of the population administered the telephone questionnaire

Gender	32 men 57 women	
Age	mean	58
	range	19-80
Visits	mean	5
	range	1-21
444 Total visits		
	pain	53%
	xerostomia	32%
	hot flashes	6%
	nausea, vomiting, anorexia	6%
	other (weight loss, smoking cessation, nonhealing wounds and so forth)	3%

varying symptoms received acupuncture in our Center’s Pain Clinic, Radiation and Medical Oncology Clinics and Breast Health Center. Most patients were referred by their physicians, but some, after learning of its availability, self-referred. Standard medical therapies were continued during acupuncture therapy, although palliative medications were discontinued if warranted. These patients had 823 visits with the single acupuncturist (RCN) during this time period. Our initial perspective was that the addition of acupuncture was a successful intervention, but prior to training more providers and further expanding services, a practice analysis was performed. It was also hoped that this would help in determining which types of complaints would be best served by acupuncture.

Techniques of acupuncture varied depending on the clinical situation. Included were traditional Chinese acupuncture, auricular acupuncture, percutaneous electrical nerve stimulation (PENS), and Korean hand and

Japanese scalp acupuncture. Disposable sterile single-use needles were used exclusively. If stimulation was required of the acupuncture point, this was accomplished with heat or electricity. In no cases was moxibustion performed. As an example, therapy for hot flashes varies between an energetic input using the Tai Yang/Shao Yin meridians and an N/N+1 circuit¹¹ with the classical points Kidney-3, Bladder-60, and Heart-3, and a classical scheme using Kidney-3, Bladder-60, Spleen-6, Stomach-36 and Large Intestine-4. Xerostomia therapy uses auricular acupuncture of three points bilaterally, and a single point on the bilateral radial index finger. Back and joint pain is generally treated with PENS, a distinctly western technique; if a depressive or anxious component is present, the patient may receive eastern regimens such as Four Gates or Internal/External Dragons. Trigger points may be deactivated by the western technique of Travell and Simons,¹² or by a modification of an eastern technique of dispersion: the Two-Needle technique.¹³

This performance analysis was performed by a single physician (GRP) with patients receiving acupuncture between 1 January 2000 and 30 April 2000. The 89 patients treated during this 4-month interval had a total of 444 visits. A simple questionnaire was administered to 79 of these patients (88.7%) available by phone during the months of June and July 2000. For ease of administration, and to impact minimally on the patients’ time, a five-question metric was used. The questionnaire is in Figure 1; demographics of the respondents are in Table 1.

It is understood that the metric and design of the questionnaire is somewhat naïve by conventional standards. It was designed, however, not to definitively document results for any individual diagnosis, but to obtain a

**PRIVATE AND CONFIDENTIAL / QUALITY ASSURANCE
Acupuncture Peer Review
Naval Medical Center San Diego**

Patient:	SSN:	Date:
Practitioner:	Reviewer:	
Informed Consent Obtained?	Yes <input type="checkbox"/>	No <input type="checkbox"/>
Needle Placement:	Yes <input type="checkbox"/>	No <input type="checkbox"/>
Duration:	Yes <input type="checkbox"/>	No <input type="checkbox"/>
Procedure:		
Electrostimulation / Frequency:	Yes <input type="checkbox"/>	No <input type="checkbox"/>
Piezostimulation:	Yes <input type="checkbox"/>	No <input type="checkbox"/>
N/A:	<input type="checkbox"/>	<input type="checkbox"/>
Assessment of Outcome:	Yes <input type="checkbox"/>	No <input type="checkbox"/>
Needle Count Correct	Yes <input type="checkbox"/>	No <input type="checkbox"/>
Patient Tolerance:	Yes <input type="checkbox"/>	No <input type="checkbox"/>
Disposition:	Yes <input type="checkbox"/>	No <input type="checkbox"/>

Figure 2 Peer review form

PRIVATE AND CONFIDENTIAL / QUALITY ASSURANCE
Acupuncture Peer Review
Naval Medical Center San Diego

May, 2001

Note documentation deficits for month:	@ ¹	#	&	*
1. Informed Consent documented				
2. Needle placement				
3. Duration	1			
4. Procedure				
5. Outcome				5
6. All needles accounted for				
7. Patient tolerance		2		
8. Disposition				
Total notes reviewed:	10	28	14	72
Total appropriate:	10	27	14	72

Notes:

- @ Legibility is an issue.
 # Consider 10 minutes per side for internal / external dragons.

¹ Identities of the individual physicians are coded for confidentiality. Each knows his/her code, and each physician signs this sheet at the monthly meeting.

Figure 3 Monthly peer review summary

broad overview of patient perspectives and results. Results for specific diagnoses are subsequently obtained in individual, in-depth surveys using specific metrics such as a visual analogue scale for pain, the Xerostomia Inventory,¹⁴ or the Mayo Clinic Hot Flash metric.¹⁵

Results

No adverse events referable to acupuncture were noted.

Of the 42 patients referred for palliation of pain, 15 (36%) had no response. However, those with a response noted a mean value of 71% improvement. When that figure is averaged over all pain patients (including those who had no response), the mean improvement was 46%. Twenty-three pain patients (55%) showed at least 30% improvement, and 14 (33%) showed at least 80% improvement. Duration of response was variable, but often resulted in lessening of oral analgesia. Further, patients stated that they had improvement in their ability to perform daily activities and tolerate the side effects from the treatment of their illness.

Of the 25 patients referred for xerostomia, 7 (28%) had no durable response. Responders experienced a mean 50% improvement, with 14 patients (56%) showing at least 30% improvement and 4 patients (16%) with at least 80% improvement. Most of these patients were previously reported using a different metric.⁵

All patients in the smaller groups referred for hot flashes, or nausea and vomiting, experienced palliation of their symptoms. The mean improvement in hot flashes was 81%, and was 96% for nausea and vomiting.

When asked the global question regarding acupuncture's worth, 68 patients (86%) considered it very important, 9 patients (11%) considered it slightly important and 2 patients (3%) considered it unimportant. Those percentages continued when patients were asked whether our centre should continue acupuncture services: 90% recommended continuing the service and 10% (8 patients) recommended discontinuation of services.

Table 2 Major diagnoses treated with acupuncture (first 2 months of expanded programme)

Diagnosis	Number of		Percentage of	
	Patients	Visits	Patients	Visits
Pain (incl.)	41	103	36%	41%
RSD	2	4		
Neuropathy	6	14		
Xerostomia	19	43	17%	17%
Hot flashes	8	26	7%	10%
Obesity	3	13	3%	5%
Protocol (IRB approved, for lower urinary tract symptoms)	8	28	7%	7%
Others	34	39		
Total	113	252		

Discussion

Success in treating xerostomia with acupuncture has been previously reported by our group.^{7,10} While we were struck by such success in the limited oncology population of patients with head and neck cancer post-radiotherapy, it was also clear that acupuncture was valuable in the larger, general oncology community. However, prior to expanding the number of acupuncture providers and offering acupuncture services to non-oncology patients, prudent business practice dictated that validation of our impressions in terms of baseline patient satisfaction was documented.

It was reproducible across the two largest diagnoses that about 30% of patients had no response to acupuncture. This is concordant with the literature.^{16,17} The cost of acupuncture is less than all but the most inexpensive pain medications (US\$0.10–US\$0.15 per sterilized, disposable acupuncture needle), and no adverse events were noted. It is somewhat time intensive for the practitioner, who happens to be a physician in our system. Non-physician acupuncturists may be integrated in the United States or elsewhere, if available – but we strongly recommend active physician oversight if this is the case.

Our current programme has four physician acupuncturists: the full-time, senior acupuncturist, and three part-time practitioners (an Anesthesiologist, a Rheumatologist and a Radiation Oncologist). Caveats of their privileges are that 100% peer review is performed of their acupuncture charts, outcome analyses are performed, and monthly meetings are held for discussion of practice issues and new data. The peer review form currently used is included in Figure 2, and the monthly summary form is in Figure 3. All patient visits are input into a database for data extraction and to provide easy recall for credentialing.

In the first 2 months of the integrated programme, over 200 patient visits occurred; the most frequent are listed by diagnosis in Table 2. Outcome analyses by diagnosis are being collated and will be submitted for publication as follow-up matures.

Conclusion

These data document that acupuncture is safe, inexpensive and is considered an important service by our patients. Our previous data have revealed its use for palliation of pilocarpine-resistant xerostomia after head and neck radiotherapy.⁷ We concur with other investigators that it is a useful adjunct in chemotherapy-induced emesis,⁹ and as an adjunct mechanism of pain control.

References

- Eisenberg DM, Kessler RC, Foster C, Norlock FE, Calkins DR, Delbanco TL. Unconventional medicine in the United States: prevalence, costs, and patterns of use. *N Engl J Med* 1993; **328**: 246–52.
- Astin JA. Why patients use alternative medicine: results of a national study. *JAMA* 1998; **279**: 1548–53.
- National Institutes of Health Consensus Conference: acupuncture. *JAMA* 1998; **280**: 1518–24.
- VanTulder MW, Cherkin DC, Berman B, Lao L, Koes BW. The effectiveness of acupuncture in the management of acute and chronic low back pain. A systematic review within the framework of the Cochrane Collaboration Back Review Group. *Spine* 1999; **24**: 1113–23.
- Ezzo J, Berman B, Hadhazy VA, Jadad AR, Lao L, Singh BB. Is acupuncture effective for the treatment of chronic pain? *Pain* 2000; **86**: 217–25.
- Jacobson JS, Workman SB, Kronenberg F. Research on complementary/alternative medicine for patients with breast cancer: a review of the biomedical literature. *J Clin Oncol* 2000; **18**: 668–83.
- Johnstone PAS, Peng YP, May BC, Inouye WS, Niemtzw RC. Acupuncture for pilocarpine-resistant xerostomia after head and neck irradiation. *Int J Radiat Oncol Biol Phys* 2001; **50**: 353–57.
- Avants SK, Margolin A, Holford TR, Kosten TR. A randomized controlled trial of auricular acupuncture for cocaine dependence. *Arch Intern Med* 2000; **160**: 2305–12.
- Shen J, Wenger N, Glaspy J, Hays RD, Albert PS, Choi C, Shekelle PG. Electroacupuncture for control of myeloablative chemotherapy-induced emesis. A randomized controlled trial. *JAMA* 2000; **284**: 2755–61.
- Niemtzow RC, Peng YP, May BC, Inouye WS, Johnstone PAS. Acupuncture technique for pilocarpine-resistant xerostomia following radiotherapy for head and neck malignancies. *Med Acupunct* 2000; **12**: 42–43.
- Helms JM. *Acupuncture energetics: a clinical approach for physicians*. Berkeley, California: Medical Acupuncture Publishers, 1995.
- Travell JG, Simons DG. *Myofascial pain and dysfunction: the trigger point manual*. Baltimore: Williams and Wilkins, 1983.
- Claraco AE. The 2-needle technique. *Med Acupunct* 2000; **12**: 17–18.
- Thomson WM, Chalmers JM, Spencer AJ, Williams SM. The Xerostomia Inventory: a multi-item approach to measuring dry mouth. *Community Dent Health* 1999; **16**: 12–17.
- Quella SK, Loprinzi CL, Barton DL, *et al*. Evaluation of soy phytoestrogens for the treatment of hot flashes in breast cancer survivors: a North Central Cancer Treatment Group trial. *J Clin Oncol* 2000; **18**: 1068–74.
- Lewith GT, Machin D. On the evaluation of the clinical effects of acupuncture. *Pain* 1983; **16**: 111–127.
- Yong D, Lim SH, Zhao CX, Cui SL, Zhang L, Lee TL. Acupuncture treatment at Ang Mo Kio Community Hospital – a report on our initial experience. *Singapore Med J* 1999; **40**: 260–64.

AUTHOR QUERIES

AUTHOR PLEASE ANSWER ALL QUERIES

1. Table 2: Please check total of values in columns 2 and 3. In the second column, 113 should be 121 and in the third column, 252 should be 270.
2. Please see 88.7% in Abstract and in Patients and Methods section (paragraph 3). Journal style requires that small percentages be rounded up or down to a whole or half percentage. Please note if 88.7% should be changed to 88.5% or 89%. Thank you.

Journal Code: PM	Jobname: pm540oa	Page: 6 of 6	Date: 2/15/2002	Time: 22:4
------------------	---------------------	--------------	-----------------	------------